

Amendments to the Claims:

The listing replaces all prior versions and listings of claims in this application.

Listing of Claims:

1. (Currently amended) A method for treating prostatic intraepithelial neoplasia in a subject, comprising the step of administering an effective amount of a composition to said subject to treat ~~or prevent~~ said prostatic intraepithelial neoplasia, said composition comprising therapeutically effective amounts of supercritical extracts of rosemary, turmeric, oregano and ginger; and therapeutically effective amounts of hydroalcoholic extracts of holy basil, ginger, turmeric, Scutellaria baicalensis, rosemary, green tea, huzhang, Chinese goldthread, and barberry.

2. (Original) The method of claim 1, wherein said composition is administered orally.

3. (Original) The method of claim 2, wherein the orally administered composition is in the form of one or more capsules, one or more tablets, or one or more pills.

4. (Original) The method of claim 1, wherein the composition comprises:

(A) from about 4.5% to about 7.5% by weight of the hydroalcoholic extract of ginger;

(B) from about 5.5% to about 8.5% by weight of the supercritical extract of ginger;

(C) from about 1.0% to about 1.5% by weight of the supercritical extract of turmeric;

(D) from about 10.0% to about 16.0% by weight of the supercritical extract of rosemary;

(E) from about 4.0% to about 6.0% by weight of the supercritical extract of oregano;

(F) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of turmeric;

(G) from about 5.5% to about 8.0% by weight of the hydroalcoholic extract of rosemary;

(H) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of holy basil;

(I) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of green tea;

(J) from about 8.0% to about 12.0% by weight of the hydroalcoholic extract of huzhang;

(K) from about 4.0% to about 6.0% by weight of the hydroalcoholic extract of Chinese goldthread;

(L) from about 4.0% to about 6.0% by weight of the

hydroalcoholic extract of barberry; and

(M) from about 2.0% to about 3.0% by weight of the hydroalcoholic extract of *Scutellaria baicalensis*.

5. (Original) The method of claim 1, wherein the weight ratio of the supercritical extract of ginger to the hydroalcoholic extract of ginger is from about 0.9:1 to about 1.4:1.

6. (Original) The method of claim 1, wherein the weight ratio of the hydroalcoholic extract of turmeric to the supercritical extract of turmeric is from about 8:1 to about 12:1.

7. (Original) The method of claim 1, wherein the weight ratio of the supercritical extract of rosemary to the hydroalcoholic extract of rosemary is from about 1.6:1 to about 2.4:1.

8. (Original) The method of claim 1, wherein the hydroalcoholic extract of ginger comprises from about 2.4% to about 3.6% by weight of pungent compounds.

9. (Original) The method of claim 1, wherein the supercritical extract of ginger comprises from about 24% to about 36% by weight

of pungent compounds and from about 6.4% to about 9.6% by weight of zingiberene.

10. (Original) The method of claim 1, wherein the supercritical extract of turmeric comprises from about 36% to about 54% by weight of turmerones.

11. (Original) The method of claim 1, wherein the supercritical extract of rosemary comprises from about 18.4% to about 27.6% by weight of total phenolic antioxidants.

12. (Original) The method of claim 1, wherein the supercritical extract of oregano comprises from about 0.64% to about 0.96% by weight of total phenolic antioxidants.

13. (Original) The method of claim 1, wherein the hydroalcoholic extract of turmeric comprises from about 5.6% to about 8.4% by weight of curcumin.

14. (Original) The method of claim 1, wherein the hydroalcoholic extract of rosemary comprises from about 18.4% to about 27.6% by weight of total phenolic antioxidants.

15. (Original) The method of claim 1, wherein the hydroalcoholic extract of holy basil comprises from about 1.6% to about 2.4% by weight of ursolic acid.

16. (Original) The method of claim 1, wherein the hydroalcoholic extract of green tea comprises from about 36% to about 54% by weight of polyphenols.

17. (Original) The method of claim 1, wherein the hydroalcoholic extract of huzhang comprises from about 6.4% to about 9.6% by weight of resveratrol.

18. (Original) The method of claim 1, wherein the hydroalcoholic extract of Chinese goldthread comprises from about 4.8% to about 7.2% by weight of berberine.

19. (Original) The method of claim 1, wherein the hydroalcoholic extract of barberry comprises from about 4.8% to about 7.2% by weight of berberine.

20. (Original) The method of claim 1, wherein said composition provided in step (a) comprises:

(A) from about 4.5% to about 7.5% by weight of the

hydroalcoholic extract of ginger, wherein the extract comprises from about 2.4% to about 3.6% by weight of pungent compounds;

(B) from about 5.5% to about 8.5% by weight of the supercritical extract of ginger, wherein the extract comprises from about 24% to about 36% by weight of pungent compounds and from about 6.4% to about 9.6% by weight of zingiberene;

(C) from about 1.0% to about 1.5% by weight of the supercritical extract of turmeric, wherein the extract comprises from about 36% to about 54% by weight of turmerones;

(D) from about 10.0% to about 16.0% by weight of the supercritical extract of rosemary, wherein the extract comprises from about 18.4% to about 27.6% by weight of total phenolic antioxidants;

(E) from about 4.0% to about 6.0% by weight of the supercritical extract of oregano, wherein the extract comprises from about 0.64% to about 0.96% by weight of total phenolic antioxidants;

(F) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of turmeric, wherein the extract comprises from about 5.6% to about 8.4% by weight of curcumin;

(G) from about 5.5% to about 8.0% by weight of the hydroalcoholic extract of rosemary, wherein the extract comprises from about 18.4% to about 27.6% by weight of total phenolic

antioxidants;

(H) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of holy basil, wherein the extract comprises from about 1.6% to about 2.4% by weight of ursolic acid;

(I) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of green tea, wherein the extract comprises from about 36% to about 54% by weight of polyphenols;

(J) from about 8.0% to about 12.0% by weight of the hydroalcoholic extract of huzhang, wherein the extract comprises from about 6.4% to about 9.6% by weight of resveratrol;

(K) from about 4.0% to about 6.0% by weight of the hydroalcoholic extract of Chinese goldthread, wherein the extract from about 4.8% to about 7.2% by weight of berberine;

(L) from about 4.0% to about 6.0% by weight of the hydroalcoholic extract of barberry, wherein the extract from about 4.8% to about 7.2% by weight of berberine; and

(M) from about 2.0% to about 3.0% by weight of the hydroalcoholic extract of *Scutellaria baicalensis*;

and wherein said composition further comprises:

(i) the supercritical extract of ginger and the hydroalcoholic extract of ginger at a weight ratio of from about 0.9 to about 1.4 parts of supercritical extract per 1 part of hydroalcoholic extract;

(ii) the hydroalcoholic extract of turmeric and the supercritical extract of turmeric at a weight ratio of from about 8 to about 12 parts of hydroalcoholic extract per 1 part of supercritical extract; and

(iii) the supercritical extract of rosemary and the hydroalcoholic extract of rosemary at a weight ratio of from about 1.6 to about 2.4 parts of supercritical extract per 1 part of hydroalcoholic extract.

21. (Original) The method of claim 1, said composition is administered in a daily dosage of at least about 700 mg.

22. (Original) The method of claim 1, wherein said composition is administered on a daily basis for at least 4 weeks.

23. (Currently amended) A method for treating precancerous cellular proliferations within prostatic ducts, ductiles and acini in a subject, comprising the step of administering an effective amount of a composition to said subject to treat ~~or prevent~~ said precancerous cellular proliferations, said composition comprising therapeutically effective amounts of supercritical extracts of rosemary, turmeric, oregano and ginger; and therapeutically

effective amounts of hydroalcoholic extracts of holy basil, ginger, turmeric, *Scutellaria baicalensis*, rosemary, green tea, huzhang, Chinese goldthread, and barberry.

24. (Original) The method of claim 23, wherein said composition is administered orally.

25. (Original) The method of claim 24, wherein the orally administered composition is in the form of one or more capsules, one or more tablets, or one or more pills.

26. (Original) The method of claim 23, wherein the composition comprises:

(A) from about 4.5% to about 7.5% by weight of the hydroalcoholic extract of ginger;

(B) from about 5.5% to about 8.5% by weight of the supercritical extract of ginger;

(C) from about 1.0% to about 1.5% by weight of the supercritical extract of turmeric;

(D) from about 10.0% to about 16.0% by weight of the supercritical extract of rosemary;

(E) from about 4.0% to about 6.0% by weight of the supercritical extract of oregano;

(F) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of turmeric;

(G) from about 5.5% to about 8.0% by weight of the hydroalcoholic extract of rosemary;

(H) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of holy basil;

(I) from about 10.0% to about 16.0% by weight of the hydroalcoholic extract of green tea;

(J) from about 8.0% to about 12.0% by weight of the hydroalcoholic extract of huzhang;

(K) from about 4.0% to about 6.0% by weight of the hydroalcoholic extract of Chinese goldthread;

(L) from about 4.0% to about 6.0% by weight of the hydroalcoholic extract of barberry; and

(M) from about 2.0% to about 3.0% by weight of the hydroalcoholic extract of *Scutellaria baicalensis*.

27. (Original) The method of claim 23, wherein the weight ratio of the supercritical extract of ginger to the hydroalcoholic extract of ginger is from about 0.9:1 to about 1.4:1.

28. (Original) The method of claim 23, wherein the weight ratio of the hydroalcoholic extract of turmeric to the

supercritical extract of turmeric is from about 8:1 to about 12:1.

29. (Original) The method of claim 23, wherein the weight ratio of the supercritical extract of rosemary to the hydroalcoholic extract of rosemary is from about 1.6:1 to about 2.4:1.

30. (Original) The method of claim 23, wherein the post supercritical hydroalcoholic extract of ginger comprises from about 2.4% to about 3.6% by weight of pungent compounds.

31. (Original) The method of claim 23, wherein the supercritical extract of ginger comprises from about 24% to about 36% by weight of pungent compounds and from about 6.4% to about 9.6% by weight of zingiberene.

32. (Original) The method of claim 23, wherein the supercritical extract of turmeric comprises from about 36% to about 54% by weight of turmerones.

33. (Original) The method of claim 23, wherein the supercritical extract of rosemary comprises from about 18.4% to about 27.6% by weight of total phenolic antioxidants.

34. (Original) The method of claim 23, wherein the supercritical extract of oregano comprises from about 0.64% to about 0.96% by weight of total phenolic antioxidants.

35. (Original) The method of claim 23, wherein the hydroalcoholic extract of turmeric comprises from about 5.6% to about 8.4% by weight of curcumin.

36. (Original) The method of claim 23, wherein the hydroalcoholic extract of rosemary comprises from about 18.4% to about 27.6% by weight of total phenolic antioxidants.

37. (Original) The method of claim 23, wherein the hydroalcoholic extract of holy basil comprises from about 1.6% to about 2.4% by weight of ursolic acid.

38. (Original) The method of claim 23, wherein the hydroalcoholic extract of green tea comprises from about 36% to about 54% by weight of polyphenols.

39. (Original) The method of claim 23, wherein the hydroalcoholic extract of huzhang comprises from about 6.4% to

about 9.6% by weight of resveratrol.

40. (Original) The method of claim 23, wherein the hydroalcoholic extract of Chinese goldthread comprises from about 4.8% to about 7.2% by weight of berberine.

41. (Original) The method of claim 23, wherein the hydroalcoholic extract of barberry comprises from about 4.8% to about 7.2% by weight of berberine.

42. (Original) The method of claim 23, said composition is administered in a daily dosage of at least about 700 mg.

43. (Original) The method of claim 23, wherein said composition is administered on a daily basis for at least 4 weeks.